SYNTHETIC EXAMPLE

AUDIT

AUDIT - synthetic example

CLIENT	ADMINISTERED	RESPONSE MODE
John Doe	**Example date**	**Self-report**

TOTAL

23 / 40

DEMO SYNTHETIC



23/40

What this report lays out

- Source instrument: AUDIT
- Item count: 10 items
- Response format: Varies by item
- Respondent: Patient

CLINICIAN REVIEW REQUIRED

This document demonstrates Lirena's report design using fictional data. It does not represent a scored AUDIT result, cannot establish a diagnosis, and must not be placed in a clinical record.

AUDIT does not diagnose alcohol use disorder or determine detox need. Those are clinical decisions.

A report designed for clinical review

Synthetic AUDIT example. The gauges and rows use this instrument's groups and range. They are not a client record.

REVIEW

Adults complete ten items. Positive findings should trigger fuller assessment and local alcohol-care pathways, including safety review where relevant.

CONTEXT

A WHO screen of alcohol consumption, dependence symptoms and alcohol-related harm for clinician review. It does not diagnose an alcohol use disorder. The implemented range is 0 to 40. Lirena's implemented cutoff is 8.

BOUNDARY

AUDIT does not diagnose alcohol use disorder or determine detox need. Those are clinical decisions.

REPORT PRINCIPLE

Show the evidence, the version, and the boundary. Leave diagnosis and formulation with the clinician.

AUDIT response pattern

Synthetic responses for AUDIT. The table uses this instrument's groups and range. It is not a client record and does not reproduce official item wording.

ITEM	GROUP	SYNTHETIC RESPONSE
1	Total	Synthetic response
2	Total	Synthetic response
3	Total	Synthetic response
4	Total	Synthetic response
5	Total	Synthetic response
6	Total	Synthetic response
7	Total	Synthetic response
8	Total	Synthetic response
9	Total	Synthetic response
10	Total	Synthetic response

Synthetic AUDIT values only. Not a client record.

Source-led by design

Lirena separates published methodology, product implementation, and clinical approval. A citation does not by itself authorize administration or prove a software scoring rule.

INSTRUMENT

The full AUDIT includes consumption items plus questions about control, morning drinking, guilt, blackouts, injury, and concern from others. It is still a screen, not a specialist dependence assessment.

ADMINISTRATION

Adults complete ten items. Positive findings should trigger fuller assessment and local alcohol-care pathways, including safety review where relevant.

CLINICAL BOUNDARY

AUDIT does not diagnose alcohol use disorder or determine detox need. Those are clinical decisions.

METHODOLOGY

A WHO screen of alcohol consumption, dependence symptoms and alcohol-related harm for clinician review. It does not diagnose an alcohol use disorder. The implemented range is 0 to 40. Lirena's implemented cutoff is 8.

SELECTED REFERENCES

1. WHO AUDIT manual
World Health Organization. AUDIT: the Alcohol Use Disorders Identification Test.

2. Scoring source Lirena follows
Source URL recorded on the Lirena scoring verification input.