SYNTHETIC EXAMPLE

C-SSRS screener

C-SSRS screener - synthetic example

CLIENT
John Doe

ADMINISTERED
Example date

RESPONSE MODE
Self-report

HIGHEST IDEATION QUESTION ANSWERED YES

4 / 5

DEMO SYNTHETIC



BEHAVIOR IN THE PAST 3 MONTHS

1 / 1

DEMO SYNTHETIC



What this report lays out

- Source instrument: C-SSRS screener
- Item count: 7 items
- Response format: Yes, No
- Respondent: Patient

CLINICIAN REVIEW REQUIRED

This document demonstrates Lirena's report design using fictional data. It does not represent a scored C-SSRS result, cannot establish a diagnosis, and must not be placed in a clinical record.

C-SSRS does not determine risk by itself, does not provide emergency care, and does not replace a clinician's assessment or local protocol. Absence of reported ideation is not a guarantee of safety.

A report designed for clinical review

Synthetic C-SSRS screener example. The gauges and rows use this instrument's groups and range. They are not a client record.

REVIEW

The stored form is the 2026 self-report recent screener, completed by the respondent and reviewed by the clinic. This page is not a crisis service, and Lirena does not monitor responses in real time.

CONTEXT

Lirena uses this rule. Score range: Highest ideation question answered Yes (05) and behavior in the past 3 months (0 or 1). No total and no severity band. The two readings follow the 2026 self-report recent form. C-SSRS screener answers are collected for clinician review. Lirena does not assign a risk category or a safety clearance. Item groups: Highest ideation question answered Yes (membership on the scoring rule, 0 to 5); Behavior in the past 3 months (membership on the scoring rule, 0 to 1). Lirena does not combine these groups into one total. Cutoff not implemented in Lirena.

BOUNDARY

C-SSRS does not determine risk by itself, does not provide emergency care, and does not replace a clinician's assessment or local protocol. Absence of reported ideation is not a guarantee of safety.

REPORT PRINCIPLE

Show the evidence, the version, and the boundary. Leave diagnosis and formulation with the clinician.

C-SSRS screener response pattern

Synthetic responses for C-SSRS screener. The table uses this instrument's groups and range. It is not a client record and does not reproduce official item wording.

ITEM	GROUP	SYNTHETIC RESPONSE
1	Highest ideation question answered Yes	**Yes**
2	Behavior in the past 3 months	**No**
3	Highest ideation question answered Yes	**Yes**
4	Behavior in the past 3 months	**No**
5	Highest ideation question answered Yes	**Yes**
6	Behavior in the past 3 months	**No**
7	Highest ideation question answered Yes	**Yes**

Synthetic C-SSRS screener values only. Not a client record.

Source-led by design

Lirena separates published methodology, product implementation, and clinical approval. A citation does not by itself authorize administration or prove a software scoring rule.

INSTRUMENT

The C-SSRS, developed at Columbia University, is a family of tools for asking about suicidal ideation and behaviour in a consistent way. This catalogue entry refers to a screener form, not every C-SSRS version. It is screening support inside a clinic-owned protocol.

ADMINISTRATION

The stored form is the 2026 self-report recent screener, completed by the respondent and reviewed by the clinic. This page is not a crisis service, and Lirena does not monitor responses in real time.

CLINICAL BOUNDARY

C-SSRS does not determine risk by itself, does not provide emergency care, and does not replace a clinician's assessment or local protocol. Absence of reported ideation is not a guarantee of safety.

METHODOLOGY

Lirena uses this rule. Score range: Highest ideation question answered Yes (05) and behavior in the past 3 months (0 or 1). No total and no severity band. The two readings follow the 2026 self-report recent form. C-SSRS screener answers are collected for clinician review. Lirena does not assign a risk category or a safety clearance. Item groups: Highest ideation question answered Yes (membership on the scoring rule, 0 to 5); Behavior in the past 3 months (membership on the scoring rule, 0 to 1). Lirena does not combine these groups into one total. Cutoff not implemented in Lirena.

SELECTED REFERENCES

1. Official C-SSRS programme
Columbia Lighthouse Project. Columbia Suicide Severity Rating Scale.

2. Scale development paper
Posner K et al. The Columbia-Suicide Severity Rating Scale. Am J Psychiatry. 2011;168(12):1266-1277.

3. Scoring source Lirena follows
Source URL recorded on the Lirena scoring verification input.