SYNTHETIC EXAMPLE

DAST-10

DAST-10 - synthetic example

CLIENT	ADMINISTERED	RESPONSE MODE
John Doe	**Example date**	**Self-report**

TOTAL

6 / 10

DEMO SYNTHETIC



6/10

What this report lays out

- Source instrument: DAST-10
- Item count: 10 items
- Response format: Yes, No
- Respondent: Patient

CLINICIAN REVIEW REQUIRED

This document demonstrates Lirena's report design using fictional data. It does not represent a scored DAST-10 result, cannot establish a diagnosis, and must not be placed in a clinical record.

DAST-10 does not diagnose a substance use disorder, identify which drug is involved, or determine treatment.

A report designed for clinical review

Synthetic DAST-10 example. The gauges and rows use this instrument's groups and range. They are not a client record.

REVIEW

Adults complete ten yes/no items. Positive screens should trigger fuller assessment, local policy, and safety review. This page does not collect those answers.

CONTEXT

A brief self-report screen of problems related to drug use other than alcohol, for clinician review. It does not diagnose a substance use disorder. The implemented range is 0 to 10. Lirena's implemented cutoff is 3.

BOUNDARY

DAST-10 does not diagnose a substance use disorder, identify which drug is involved, or determine treatment.

REPORT PRINCIPLE

Show the evidence, the version, and the boundary. Leave diagnosis and formulation with the clinician.

DAST-10 response pattern

Synthetic responses for DAST-10. The table uses this instrument's groups and range. It is not a client record and does not reproduce official item wording.

ITEM	GROUP	SYNTHETIC RESPONSE
1	Total	Yes
2	Total	No
3	Total	Yes
4	Total	No
5	Total	Yes
6	Total	No
7	Total	Yes
8	Total	No
9	Total	Yes
10	Total	No

Synthetic DAST-10 values only. Not a client record.

Source-led by design

Lirena separates published methodology, product implementation, and clinical approval. A citation does not by itself authorize administration or prove a software scoring rule.

INSTRUMENT

The Drug Abuse Screening Test-10 asks about problems related to drugs other than alcohol over the past 12 months. It is a brief screen, not a toxicology result and not a full substance-use interview.

ADMINISTRATION

Adults complete ten yes/no items. Positive screens should trigger fuller assessment, local policy, and safety review. This page does not collect those answers.

CLINICAL BOUNDARY

DAST-10 does not diagnose a substance use disorder, identify which drug is involved, or determine treatment.

METHODOLOGY

A brief self-report screen of problems related to drug use other than alcohol, for clinician review. It does not diagnose a substance use disorder. The implemented range is 0 to 10. Lirena's implemented cutoff is 3.

SELECTED REFERENCES

1. DAST development
Skinner HA. The drug abuse screening test. Addict Behav. 1982;7(4):363-371.

2. Scoring source Lirena follows
Source URL recorded on the Lirena scoring verification input.

3. Developer or publisher page
CAMH DAST information