SYNTHETIC EXAMPLE

DS-II

DS-II - synthetic example

CLIENT
John Doe

ADMINISTERED
Example date

RESPONSE MODE
Self-report

TOTAL

28 / 32

DEMO SYNTHETIC



What this report lays out

- Source instrument: DS-II
- Item count: 16 items
- Response format: Never, Sometimes, Often
- Respondent: Self-report

CLINICIAN REVIEW REQUIRED

This document demonstrates Lirena's report design using fictional data. It does not represent a scored DS-II result, cannot establish a diagnosis, and must not be placed in a clinical record.

DS-II is not a diagnosis. A result cannot replace clinician-led assessment, history, examination, or investigation where those are required.

A report designed for clinical review

Synthetic DS-II example. The gauges and rows use this instrument's groups and range. They are not a client record.

REVIEW

DS-II is completed by Self-report. Response options: Never, Sometimes, Often. Reference period not sourced. The result is for clinician review.

CONTEXT

Sixteen-item DS-II total from 0 to 32 with two eight-item subscales, Meaning and Purpose and Distress and Coping Ability. The bands are the published low, moderate and high ranges for clinician review.

BOUNDARY

DS-II is not a diagnosis. A result cannot replace clinician-led assessment, history, examination, or investigation where those are required.

REPORT PRINCIPLE

Show the evidence, the version, and the boundary. Leave diagnosis and formulation with the clinician.

DS-II response pattern

Synthetic responses for DS-II. The table uses this instrument's groups and range. It is not a client record and does not reproduce official item wording.

ITEM	GROUP	SYNTHETIC RESPONSE
1	Total	Never
2	Total	Sometimes
3	Total	Often
4	Total	Never
5	Total	Sometimes
6	Total	Often
7	Total	Never
8	Total	Sometimes
9	Total	Often
10	Total	Never
11	Total	Sometimes
12	Total	Often
13	Total	Never
14	Total	Sometimes
15	Total	Often
16	Total	Never

Synthetic DS-II values only. Not a client record.

Source-led by design

Lirena separates published methodology, product implementation, and clinical approval. A citation does not by itself authorize administration or prove a software scoring rule.

INSTRUMENT

The Demoralization Scale-II asks about meaning, purpose, and coping over the last two weeks. Its published low (0 to 3), moderate (4 to 10) and high (11 or more) ranges are prompts for clinician review. It is not a diagnosis and not an end-of-life plan.

ADMINISTRATION

DS-II is completed by Self-report. Response options: Never, Sometimes, Often. Reference period not sourced. The result is for clinician review.

CLINICAL BOUNDARY

DS-II is not a diagnosis. A result cannot replace clinician-led assessment, history, examination, or investigation where those are required.

METHODOLOGY

Sixteen-item DS-II total from 0 to 32 with two eight-item subscales, Meaning and Purpose and Distress and Coping Ability. The bands are the published low, moderate and high ranges for clinician review.

SELECTED REFERENCES

1. DS-II source literature

Robinson S, Kissane DW, Brooker J, et al. Refinement and revalidation of the demoralization scale: the DS-II-internal validity. Cancer. 2016;122(14):2251-2259.

2. Scoring source Lirena follows

Source URL recorded on the Lirena scoring verification input.