SYNTHETIC EXAMPLE

EPDS

EPDS - synthetic example

CLIENT
John Doe

ADMINISTERED
Example date

RESPONSE MODE
Self-report

TOTAL

27 / 30

DEMO SYNTHETIC



27/30

What this report lays out

- Source instrument: EPDS
- Item count: 10 items
- Reference period: Past 7 days
- Response format: Varies by item

CLINICIAN REVIEW REQUIRED

This document demonstrates Lirena's report design using fictional data. It does not represent a scored EPDS result, cannot establish a diagnosis, and must not be placed in a clinical record.

EPDS does not diagnose postnatal depression, anxiety, or psychosis. A low score does not rule out illness. Any concerning item requires clinician-led review.

A report designed for clinical review

Synthetic EPDS example. The gauges and rows use this instrument's groups and range. They are not a client record.

REVIEW

The person completes ten items about the past seven days. Results should reach a clinician who understands perinatal context, social support, and local safeguarding. Lirena does not provide urgent monitoring.

CONTEXT

The EPDS is a screening measure, not a diagnosis. Any response other than Never to item 10 requires direct clinical review. The implemented range is 0 to 30.

BOUNDARY

EPDS does not diagnose postnatal depression, anxiety, or psychosis. A low score does not rule out illness. Any concerning item requires clinician-led review.

REPORT PRINCIPLE

Show the evidence, the version, and the boundary. Leave diagnosis and formulation with the clinician.

EPDS response pattern

Synthetic responses for EPDS. The table uses this instrument's groups and range. It is not a client record and does not reproduce official item wording.

ITEM	GROUP	SYNTHETIC RESPONSE
1	Total	Synthetic response
2	Total	Synthetic response
3	Total	Synthetic response
4	Total	Synthetic response
5	Total	Synthetic response
6	Total	Synthetic response
7	Total	Synthetic response
8	Total	Synthetic response
9	Total	Synthetic response
10	Total	Synthetic response

Synthetic EPDS values only. Not a client record.

Source-led by design

Lirena separates published methodology, product implementation, and clinical approval. A citation does not by itself authorize administration or prove a software scoring rule.

INSTRUMENT

The EPDS asks about how a pregnant or postnatal person has felt in the past week. It is widely used as a perinatal screening aid. Item-level review, including any self-harm-related response, remains a clinician responsibility.

ADMINISTRATION

The person completes ten items about the past seven days. Results should reach a clinician who understands perinatal context, social support, and local safeguarding. Lirena does not provide urgent monitoring.

CLINICAL BOUNDARY

EPDS does not diagnose postnatal depression, anxiety, or psychosis. A low score does not rule out illness. Any concerning item requires clinician-led review.

METHODOLOGY

The EPDS is a screening measure, not a diagnosis. Any response other than Never to item 10 requires direct clinical review. The implemented range is 0 to 30.

SELECTED REFERENCES

1. Original EPDS paper

Cox JL, Holden JM, Sagovsky R. Detection of postnatal depression. Br J Psychiatry. 1987;150:782-786.

2. Scoring source Lirena follows

Source URL recorded on the Lirena scoring verification input.