SYNTHETIC EXAMPLE

FCRI-SF

FCRI-SF - synthetic example

CLIENT
John Doe

ADMINISTERED
Example date

RESPONSE MODE
Self-report

TOTAL

23 / 36

DEMO SYNTHETIC



23/36

What this report lays out

- Source instrument: FCRI-SF
- Item count: 9 items
- Response format: Varies by item
- Respondent: Self-report

CLINICIAN REVIEW REQUIRED

This document demonstrates Lirena's report design using fictional data. It does not represent a scored FCRI-SF result, cannot establish a diagnosis, and must not be placed in a clinical record.

FCRI-SF is not a diagnosis. A result cannot replace clinician-led assessment, history, examination, or investigation where those are required.

A report designed for clinical review

Synthetic FCRI-SF example. The gauges and rows use this instrument's groups and range. They are not a client record.

REVIEW

FCRI-SF is completed by Self-report. Response format varies by item. Reference period not sourced. The result is for clinician review.

CONTEXT

Nine-item FCRI-SF severity total from 0 to 36, with item 5 reverse scored. The bands are the published screening cut-off of 13 and clinical cut-off of 22 for clinician review. The total is not a diagnosis and does not predict recurrence.

BOUNDARY

FCRI-SF is not a diagnosis. A result cannot replace clinician-led assessment, history, examination, or investigation where those are required.

REPORT PRINCIPLE

Show the evidence, the version, and the boundary. Leave diagnosis and formulation with the clinician.

FCRI-SF response pattern

Synthetic responses for FCRI-SF. The table uses this instrument's groups and range. It is not a client record and does not reproduce official item wording.

ITEM	GROUP	SYNTHETIC RESPONSE
1	Total	Synthetic response
2	Total	Synthetic response
3	Total	Synthetic response
4	Total	Synthetic response
5	Total	Synthetic response
6	Total	Synthetic response
7	Total	Synthetic response
8	Total	Synthetic response
9	Total	Synthetic response

Synthetic FCRI-SF values only. Not a client record.

Source-led by design

Lirena separates published methodology, product implementation, and clinical approval. A citation does not by itself authorize administration or prove a software scoring rule.

INSTRUMENT

The Fear of Cancer Recurrence Inventory Short Form organises how strongly a person worries that cancer will return. Published cut-offs of 13 for screening and 22 for clinical fear of recurrence are prompts for clinician review. The total is not a diagnosis and not a treatment recommendation.

ADMINISTRATION

FCRI-SF is completed by Self-report. Response format varies by item. Reference period not sourced. The result is for clinician review.

CLINICAL BOUNDARY

FCRI-SF is not a diagnosis. A result cannot replace clinician-led assessment, history, examination, or investigation where those are required.

METHODOLOGY

Nine-item FCRI-SF severity total from 0 to 36, with item 5 reverse scored. The bands are the published screening cut-off of 13 and clinical cut-off of 22 for clinician review. The total is not a diagnosis and does not predict recurrence.

SELECTED REFERENCES

1. FCRI-SF source literature

Simard S, Savard J. Fear of Cancer Recurrence Inventory: development and initial validation of a multidimensional measure of fear of cancer recurrence. Support Care Cancer. 2009;17(3):241-251.

2. FCRI-SF form

Simard S, Savard J. Fear of Cancer Recurrence Inventory - Short Form (FCRI-SF), 2015. CFPlus supplement to Can Fam Physician. 2020;66(9):672.

3. Scoring source Lirena follows

Source URL recorded on the Lirena scoring verification input.