SYNTHETIC EXAMPLE

FOSQ

FOSQ - synthetic example

CLIENT
John Doe

ADMINISTERED
Example date

RESPONSE MODE
Self-report

GENERAL PRODUCTIVITY

2 / 4

DEMO SYNTHETIC



SOCIAL OUTCOME

2 / 4

DEMO SYNTHETIC



ACTIVITY LEVEL

2 / 4

DEMO SYNTHETIC



What this report lays out

- Source instrument: FOSQ
- Item count: 31 items
- Response format: Varies by item
- Respondent: Patient

CLINICIAN REVIEW REQUIRED

This document demonstrates Lirena's report design using fictional data. It does not represent a scored FOSQ result, cannot establish a diagnosis, and must not be placed in a clinical record.

FOSQ reports subscale means and their total. It does not diagnose a sleep disorder.

A report designed for clinical review

Synthetic FOSQ example. The gauges and rows use this instrument's groups and range. They are not a client record.

REVIEW

The respondent answers the 30 September 1996 items. A response of 0 means the activity is skipped for another reason, except on the two items that have no 0. An optional note about relationships is stored and not scored.

CONTEXT

Lirena uses this rule. Score range: 0.0020.00. Five subscale means (14). A subscale with no answer from 1 to 4 is 0.00 and is not added. The total is the sum of the subscale scores. The relationship note is not scored. FOSQ reports subscale means and their total. It does not diagnose a sleep disorder. Item groups: General Productivity (membership on the scoring rule, 0.00 to 4.00); Social Outcome (membership on the scoring rule, 0.00 to 4.00); Activity Level (membership on the scoring rule, 0.00 to 4.00); Vigilance (membership on the scoring rule, 0.00 to 4.00); Intimate Relationships and Sexual Activity (membership on the scoring rule, 0.00 to 4.00). 1 item is recorded for clinician review and not included in the total. Cutoff not implemented in Lirena.

BOUNDARY

FOSQ reports subscale means and their total. It does not diagnose a sleep disorder.

REPORT PRINCIPLE

Show the evidence, the version, and the boundary. Leave diagnosis and formulation with the clinician.

FOSQ response pattern

Synthetic responses for FOSQ. The table uses this instrument's groups and range. It is not a client record and does not reproduce official item wording.

ITEM	GROUP	SYNTHETIC RESPONSE
1	General Productivity	Synthetic response
2	Social Outcome	Synthetic response
3	Activity Level	Synthetic response
4	General Productivity	Synthetic response
5	Social Outcome	Synthetic response
6	Activity Level	Synthetic response
7	General Productivity	Synthetic response
8	Social Outcome	Synthetic response
9	Activity Level	Synthetic response
10	General Productivity	Synthetic response
11	Social Outcome	Synthetic response
12	Activity Level	Synthetic response
13	General Productivity	Synthetic response
14	Social Outcome	Synthetic response
15	Activity Level	Synthetic response
16	General Productivity	Synthetic response

Synthetic FOSQ values only. Not a client record.

Source-led by design

Lirena separates published methodology, product implementation, and clinical approval. A citation does not by itself authorize administration or prove a software scoring rule.

INSTRUMENT

FOSQ focuses on functional impact rather than a single sleepiness total. Clinics use it to organise sleep-related function for clinician review, not as a stand-alone sleep-disorder diagnosis.

ADMINISTRATION

The respondent answers the 30 September 1996 items. A response of 0 means the activity is skipped for another reason, except on the two items that have no 0. An optional note about relationships is stored and not scored.

CLINICAL BOUNDARY

FOSQ reports subscale means and their total. It does not diagnose a sleep disorder.

METHODOLOGY

Lirena uses this rule. Score range: 0.0020.00. Five subscale means (14). A subscale with no answer from 1 to 4 is 0.00 and is not added. The total is the sum of the subscale scores. The relationship note is not scored. FOSQ reports subscale means and their total. It does not diagnose a sleep disorder. Item groups: General Productivity (membership on the scoring rule, 0.00 to 4.00); Social Outcome (membership on the scoring rule, 0.00 to 4.00); Activity Level (membership on the scoring rule, 0.00 to 4.00); Vigilance (membership on the scoring rule, 0.00 to 4.00); Intimate Relationships and Sexual Activity (membership on the scoring rule, 0.00 to 4.00). 1 item is recorded for clinician review and not included in the total. Cutoff not implemented in Lirena.

SELECTED REFERENCES

1. Original FOSQ paper

Weaver TE et al. An instrument to measure functional status outcomes for disorders of excessive sleepiness. Sleep. 1997;20(10):835-843.

2. Scoring source Lirena follows

Source URL recorded on the Lirena scoring verification input.