SYNTHETIC EXAMPLE

IRLS

IRLS - synthetic example

CLIENT
John Doe

ADMINISTERED
Example date

RESPONSE MODE
Clinician-led interview

TOTAL

23 / 40

DEMO SYNTHETIC



23/40

What this report lays out

- Source instrument: IRLS
- Item count: 10 items
- Response format: Varies by item
- Respondent: Patient or clinician

CLINICIAN REVIEW REQUIRED

This document demonstrates Lirena's report design using fictional data. It does not represent a scored IRLS result, cannot establish a diagnosis, and must not be placed in a clinical record.

The clinician records the patient's own ratings. The IRLS rates severity; it does not diagnose restless legs syndrome.

A report designed for clinical review

Synthetic IRLS example. The gauges and rows use this instrument's groups and range. They are not a client record.

REVIEW

An examiner administers the ten questions to a patient who meets the IRLSSG diagnostic criteria for restless legs syndrome. The patient makes every rating from 0 to 4, and the examiner is available to clarify the questions. On Lirena the clinician records the patient's ratings in the workspace; the form is never sent to the client.

CONTEXT

The IRLS total is the sum of the ten patient ratings, 0 to 40, read against the published severity bands. Item 3 counts toward the total but belongs to neither subscale. The result supports clinician review and is not a diagnosis.

BOUNDARY

The clinician records the patient's own ratings. The IRLS rates severity; it does not diagnose restless legs syndrome.

REPORT PRINCIPLE

Show the evidence, the version, and the boundary. Leave diagnosis and formulation with the clinician.

IRLS response pattern

Synthetic responses for IRLS. The table uses this instrument's groups and range. It is not a client record and does not reproduce official item wording.

ITEM	GROUP	SYNTHETIC RESPONSE
1	Total	Synthetic response
2	Total	Synthetic response
3	Total	Synthetic response
4	Total	Synthetic response
5	Total	Synthetic response
6	Total	Synthetic response
7	Total	Synthetic response
8	Total	Synthetic response
9	Total	Synthetic response
10	Total	Synthetic response

Synthetic IRLS values only. Not a client record.

Source-led by design

Lirena separates published methodology, product implementation, and clinical approval. A citation does not by itself authorize administration or prove a software scoring rule.

INSTRUMENT

The IRLS organises restless-legs severity for clinician review. A score is not a stand-alone RLS diagnosis and does not prescribe medication.

ADMINISTRATION

An examiner administers the ten questions to a patient who meets the IRLSSG diagnostic criteria for restless legs syndrome. The patient makes every rating from 0 to 4, and the examiner is available to clarify the questions. On Lirena the clinician records the patient's ratings in the workspace; the form is never sent to the client.

CLINICAL BOUNDARY

The clinician records the patient's own ratings. The IRLS rates severity; it does not diagnose restless legs syndrome.

METHODOLOGY

The IRLS total is the sum of the ten patient ratings, 0 to 40, read against the published severity bands. Item 3 counts toward the total but belongs to neither subscale. The result supports clinician review and is not a diagnosis.

SELECTED REFERENCES

1. IRLS validation

Walters AS et al. Validation of the International Restless Legs Syndrome Study Group rating scale. Sleep Medicine. 2003;4(2):121-132.

2. Scoring source Lirena follows

Source URL recorded on the Lirena scoring verification input.