SYNTHETIC EXAMPLE

MADRS

MADRS - synthetic example

CLIENT
John Doe

ADMINISTERED
Example date

RESPONSE MODE
Clinician-rated

TOTAL

55 / 60

DEMO SYNTHETIC



What this report lays out

- Source instrument: MADRS
- Item count: 10 items
- Response format: Varies by item
- Respondent: Patient, clinician rated

CLINICIAN REVIEW REQUIRED

This document demonstrates Lirena's report design using fictional data. It does not represent a scored MADRS result, cannot establish a diagnosis, and must not be placed in a clinical record.

A MADRS total is a clinician's rated severity estimate, not a diagnosis and not a treatment recommendation. Interview quality, rater training, and clinical context all matter.

A report designed for clinical review

Synthetic MADRS example. The gauges and rows use this instrument's groups and range. They are not a client record.

REVIEW

A clinician interviews the person and rates each item against published anchors. This is not a client-only form. Digital workflows must preserve the rater's role and the official rating instructions.

CONTEXT

MADRS is a clinician-rated severity measure, not a diagnosis. The total is the sum of the 10 ratings. The implemented range is 0 to 60.

BOUNDARY

A MADRS total is a clinician's rated severity estimate, not a diagnosis and not a treatment recommendation. Interview quality, rater training, and clinical context all matter.

REPORT PRINCIPLE

Show the evidence, the version, and the boundary. Leave diagnosis and formulation with the clinician.

MADRS response pattern

Synthetic responses for MADRS. The table uses this instrument's groups and range. It is not a client record and does not reproduce official item wording.

ITEM	GROUP	SYNTHETIC RESPONSE
1	Total	Synthetic response
2	Total	Synthetic response
3	Total	Synthetic response
4	Total	Synthetic response
5	Total	Synthetic response
6	Total	Synthetic response
7	Total	Synthetic response
8	Total	Synthetic response
9	Total	Synthetic response
10	Total	Synthetic response

Synthetic MADRS values only. Not a client record.

Source-led by design

Lirena separates published methodology, product implementation, and clinical approval. A citation does not by itself authorize administration or prove a software scoring rule.

INSTRUMENT

MADRS was designed to be sensitive to change in depression treatment studies. A trained clinician rates defined items after interview. It is not a patient-completed questionnaire and should not be presented as one.

ADMINISTRATION

A clinician interviews the person and rates each item against published anchors. This is not a client-only form. Digital workflows must preserve the rater's role and the official rating instructions.

CLINICAL BOUNDARY

A MADRS total is a clinician's rated severity estimate, not a diagnosis and not a treatment recommendation. Interview quality, rater training, and clinical context all matter.

METHODOLOGY

MADRS is a clinician-rated severity measure, not a diagnosis. The total is the sum of the 10 ratings. The implemented range is 0 to 60.

SELECTED REFERENCES

1. Original MADRS paper

Montgomery SA, Asberg M. A new depression scale designed to be sensitive to change. Br J Psychiatry. 1979;134:382-389.

2. MADRS severity grades

Snaith RP, Harrop FM, Newby DA, Teale C. Grade scores of the Montgomery-Asberg Depression and the Clinical Anxiety Scales. Br J Psychiatry. 1986;148:599-601.

3. Scoring source Lirena follows

Source URL recorded on the Lirena scoring verification input.