SYNTHETIC EXAMPLE

MRS

MRS - synthetic example

CLIENT	ADMINISTERED	RESPONSE MODE
John Doe	**Example date**	**Self-report**

TOTAL

25 / 44

DEMO SYNTHETIC



25/44

What this report lays out

- Source instrument: MRS
- Item count: 11 items
- Response format: none, mild, moderate, severe, very severe
- Respondent: Patient

CLINICIAN REVIEW REQUIRED

This document demonstrates Lirena's report design using fictional data. It does not represent a scored MRS result, cannot establish a diagnosis, and must not be placed in a clinical record.

MRS is a symptom rating scale, not a diagnosis. The total is the sum of the psychological, somato-vegetative and urogenital subscale scores.

A report designed for clinical review

Synthetic MRS example. The gauges and rows use this instrument's groups and range. They are not a client record.

REVIEW

MRS is completed by Patient. Response options: none, mild, moderate, severe, very severe. Reference period not sourced. The result is for clinician review.

CONTEXT

The MRS total is the sum of the psychological, somato-vegetative and urogenital subscale scores. Total-score severity categories are those published by Heinemann et al. (2004). A symptom rating scale, not a diagnosis. The implemented range is 0 to 44.

BOUNDARY

MRS is a symptom rating scale, not a diagnosis. The total is the sum of the psychological, somato-vegetative and urogenital subscale scores.

REPORT PRINCIPLE

Show the evidence, the version, and the boundary. Leave diagnosis and formulation with the clinician.

MRS response pattern

Synthetic responses for MRS. The table uses this instrument's groups and range. It is not a client record and does not reproduce official item wording.

ITEM	GROUP	SYNTHETIC RESPONSE
1	Total	none
2	Total	mild
3	Total	moderate
4	Total	severe
5	Total	very severe
6	Total	none
7	Total	mild
8	Total	moderate
9	Total	severe
10	Total	very severe
11	Total	none

Synthetic MRS values only. Not a client record.

Source-led by design

Lirena separates published methodology, product implementation, and clinical approval. A citation does not by itself authorize administration or prove a software scoring rule.

INSTRUMENT

MRS asks which of 11 menopausal symptoms apply at this time and how severe each one is. It supports clinician review of psychological, somato-vegetative and urogenital symptom severity. It does not establish menopause status by itself and does not prescribe treatment.

ADMINISTRATION

MRS is completed by Patient. Response options: none, mild, moderate, severe, very severe. Reference period not sourced. The result is for clinician review.

CLINICAL BOUNDARY

MRS is a symptom rating scale, not a diagnosis. The total is the sum of the psychological, somato-vegetative and urogenital subscale scores.

METHODOLOGY

The MRS total is the sum of the psychological, somato-vegetative and urogenital subscale scores. Total-score severity categories are those published by Heinemann et al. (2004). A symptom rating scale, not a diagnosis. The implemented range is 0 to 44.

SELECTED REFERENCES

1. MRS paper
Heinemann K et al. The Menopause Rating Scale (MRS) scale: a methodological review. Health Qual Life Outcomes. 2004;2:45.

2. International versions of the MRS
Heinemann LAJ et al. International versions of the Menopause Rating Scale (MRS). Health Qual Life Outcomes. 2003;1:28.

3. Scoring source Lirena follows
Source URL recorded on the Lirena scoring verification input.