SYNTHETIC EXAMPLE

OCI-R

OCI-R - synthetic example

CLIENT
John Doe

ADMINISTERED
Example date

RESPONSE MODE
Self-report

TOTAL

43 / 72

DEMO SYNTHETIC



What this report lays out

- Source instrument: OCI-R
- Item count: 18 items
- Response format: Not at all, A little, Moderately, A lot, Extremely
- Respondent: Patient

CLINICIAN REVIEW REQUIRED

This document demonstrates Lirena's report design using fictional data. It does not represent a scored OCI-R result, cannot establish a diagnosis, and must not be placed in a clinical record.

OCI-R does not diagnose OCD or distinguish OCD from OCPD, rumination, or tics without clinician assessment.

A report designed for clinical review

Synthetic OCI-R example. The gauges and rows use this instrument's groups and range. They are not a client record.

REVIEW

Adults rate how much statements distressed or bothered them. Results support clinician review of dimension patterns, not an automated OCD label.

CONTEXT

The OCI-R total is the sum of the 18 distress ratings and each subscale is the sum of its three items. The score does not establish a diagnosis. The implemented range is 0 to 72. Lirena's implemented cutoff is 21.

BOUNDARY

OCI-R does not diagnose OCD or distinguish OCD from OCPD, rumination, or tics without clinician assessment.

REPORT PRINCIPLE

Show the evidence, the version, and the boundary. Leave diagnosis and formulation with the clinician.

OCI-R response pattern

Synthetic responses for OCI-R. The table uses this instrument's groups and range. It is not a client record and does not reproduce official item wording.

ITEM	GROUP	SYNTHETIC RESPONSE
1	Total	Not at all
2	Total	A little
3	Total	Moderately
4	Total	A lot
5	Total	Extremely
6	Total	Not at all
7	Total	A little
8	Total	Moderately
9	Total	A lot
10	Total	Extremely
11	Total	Not at all
12	Total	A little
13	Total	Moderately
14	Total	A lot
15	Total	Extremely
16	Total	Not at all

Synthetic OCI-R values only. Not a client record.

Source-led by design

Lirena separates published methodology, product implementation, and clinical approval. A citation does not by itself authorize administration or prove a software scoring rule.

INSTRUMENT

The Obsessive-Compulsive Inventory-Revised asks about several OCD-related symptom dimensions such as washing, checking, and ordering. It is a self-report inventory for screening or monitoring, not a diagnostic interview.

ADMINISTRATION

Adults rate how much statements distressed or bothered them. Results support clinician review of dimension patterns, not an automated OCD label.

CLINICAL BOUNDARY

OCI-R does not diagnose OCD or distinguish OCD from OCPD, rumination, or tics without clinician assessment.

METHODOLOGY

The OCI-R total is the sum of the 18 distress ratings and each subscale is the sum of its three items. The score does not establish a diagnosis. The implemented range is 0 to 72. Lirena's implemented cutoff is 21.

SELECTED REFERENCES

1. OCI-R paper

Foa EB et al. The Obsessive-Compulsive Inventory: development and validation of a short version. Psychol Assess. 2002;14(4):485-496.

2. Scoring source Lirena follows

Source URL recorded on the Lirena scoring verification input.