SYNTHETIC EXAMPLE

painDETECT

painDETECT - synthetic example

CLIENT
John Doe

ADMINISTERED
Example date

RESPONSE MODE
Self-report

FINAL SCORE

35 / 38

DEMO SYNTHETIC



35/38

What this report lays out

- Source instrument: painDETECT
- Item count: 12 items
- Response format: Varies by item
- Respondent: Patient

CLINICIAN REVIEW REQUIRED

This document demonstrates Lirena's report design using fictional data. It does not represent a scored PainDetect result, cannot establish a diagnosis, and must not be placed in a clinical record.

A painDETECT band is a screening range. It is not a diagnosis of neuropathic pain and does not replace examination.

A report designed for clinical review

Synthetic painDETECT example. The gauges and rows use this instrument's groups and range. They are not a client record.

REVIEW

painDETECT is completed by Patient. Response format varies by item. Reference period not sourced. The result is for clinician review.

CONTEXT

Lirena uses this rule. Score range: Final score 038. Nociceptive (012), unclear (1318), neuropathic (1938). The published sheet also describes a score of -1; that pattern is reported as 0 and stays nociceptive. A painDETECT band is a screening range. It is not a diagnosis of neuropathic pain and does not replace examination. 3 items are recorded for clinician review and not included in the total.

BOUNDARY

A painDETECT band is a screening range. It is not a diagnosis of neuropathic pain and does not replace examination.

REPORT PRINCIPLE

Show the evidence, the version, and the boundary. Leave diagnosis and formulation with the clinician.

painDETECT response pattern

Synthetic responses for painDETECT. The table uses this instrument's groups and range. It is not a client record and does not reproduce official item wording.

ITEM	GROUP	SYNTHETIC RESPONSE
1	Final score	Synthetic response
2	Final score	Synthetic response
3	Final score	Synthetic response
4	Final score	Synthetic response
5	Final score	Synthetic response
6	Final score	Synthetic response
7	Final score	Synthetic response
8	Final score	Synthetic response
9	Final score	Synthetic response
10	Final score	Synthetic response
11	Final score	Synthetic response
12	Final score	Synthetic response

Synthetic painDETECT values only. Not a client record.

Source-led by design

Lirena separates published methodology, product implementation, and clinical approval. A citation does not by itself authorize administration or prove a software scoring rule.

INSTRUMENT

painDETECT organises symptom features that clinics may review when considering neuropathic contributions to pain. It is not interchangeable with DN4 or LANSS.

ADMINISTRATION

painDETECT is completed by Patient. Response format varies by item. Reference period not sourced. The result is for clinician review.

CLINICAL BOUNDARY

A painDETECT band is a screening range. It is not a diagnosis of neuropathic pain and does not replace examination.

METHODOLOGY

Lirena uses this rule. Score range: Final score 038. Nociceptive (012), unclear (1318), neuropathic (1938). The published sheet also describes a score of -1; that pattern is reported as 0 and stays nociceptive. A painDETECT band is a screening range. It is not a diagnosis of neuropathic pain and does not replace examination. 3 items are recorded for clinician review and not included in the total.

SELECTED REFERENCES

1. Original painDETECT paper

Freynhagen R et al. painDETECT: a new screening questionnaire to identify neuropathic components in patients with back pain. Curr Med Res Opin. 2006;22(10):1911-1920.

2. Scoring source Lirena follows

Source URL recorded on the Lirena scoring verification input.