SYNTHETIC EXAMPLE

PANSS

PANSS - synthetic example

CLIENT
John Doe

ADMINISTERED
Example date

RESPONSE MODE
Clinician-rated

TOTAL

46 / 210

DEMO SYNTHETIC



46/210

What this report lays out

- Source instrument: PANSS
- Item count: 30 items
- Response format: 7 options
- Respondent: Clinician

CLINICIAN REVIEW REQUIRED

This document demonstrates Lirena's report design using fictional data. It does not represent a scored PANSS result, cannot establish a diagnosis, and must not be placed in a clinical record.

PANSS does not diagnose schizophrenia. It is a specialist severity rating inside clinician-led care.

A report designed for clinical review

Synthetic PANSS example. The gauges and rows use this instrument's groups and range. They are not a client record.

REVIEW

A trained rater interviews the person and records the 30 ratings in the clinician completion view. Lirena does not send PANSS to the client.

CONTEXT

The PANSS total and the positive, negative and general psychopathology scale scores are raw sums of 1-7 ratings (Kay, Fiszbein and Opler, 1987). No cutoff is applied. It is a clinician rating, not a diagnosis. The implemented range is 30 to 210. Cutoff not implemented in Lirena.

BOUNDARY

PANSS does not diagnose schizophrenia. It is a specialist severity rating inside clinician-led care.

REPORT PRINCIPLE

Show the evidence, the version, and the boundary. Leave diagnosis and formulation with the clinician.

PANSS response pattern

Synthetic responses for PANSS. The table uses this instrument's groups and range. It is not a client record and does not reproduce official item wording.

ITEM	GROUP	SYNTHETIC RESPONSE
1	Total	Absent
2	Total	Minimal
3	Total	Mild
4	Total	Moderate
5	Total	Moderate severe
6	Total	Severe
7	Total	Extreme
8	Total	Absent
9	Total	Minimal
10	Total	Mild
11	Total	Moderate
12	Total	Moderate severe
13	Total	Severe
14	Total	Extreme
15	Total	Absent
16	Total	Minimal

Synthetic PANSS values only. Not a client record.

Source-led by design

Lirena separates published methodology, product implementation, and clinical approval. A citation does not by itself authorize administration or prove a software scoring rule.

INSTRUMENT

The Positive and Negative Syndrome Scale is used in specialist psychosis services and trials to rate positive, negative, and general psychopathology items after interview and observation.

ADMINISTRATION

A trained rater interviews the person and records the 30 ratings in the clinician completion view. Lirena does not send PANSS to the client.

CLINICAL BOUNDARY

PANSS does not diagnose schizophrenia. It is a specialist severity rating inside clinician-led care.

METHODOLOGY

The PANSS total and the positive, negative and general psychopathology scale scores are raw sums of 1-7 ratings (Kay, Fiszbein and Opler, 1987). No cutoff is applied. It is a clinician rating, not a diagnosis. The implemented range is 30 to 210. Cutoff not implemented in Lirena.

SELECTED REFERENCES

1. Original PANSS paper
Kay SR, Fiszbein A, Opler LA. The positive and negative syndrome scale (PANSS). Schizophr Bull. 1987;13(2):261-276.

2. Scoring source Lirena follows
Source URL recorded on the Lirena scoring verification input.

3. Developer or publisher page
Publisher PANSS information