SYNTHETIC EXAMPLE

PSEQ

PSEQ - synthetic example

CLIENT
John Doe

ADMINISTERED
Example date

RESPONSE MODE
Self-report

TOTAL

55 / 60

DEMO SYNTHETIC



What this report lays out

- Source instrument: PSEQ
- Item count: 10 items
- Response format: 7 options
- Respondent: Patient

CLINICIAN REVIEW REQUIRED

This document demonstrates Lirena's report design using fictional data. It does not represent a scored PSEQ result, cannot establish a diagnosis, and must not be placed in a clinical record.

PSEQ is not a diagnosis. A result cannot replace clinician-led assessment, history, examination, or investigation where those are required.

A report designed for clinical review

Synthetic PSEQ example. The gauges and rows use this instrument's groups and range. They are not a client record.

REVIEW

PSEQ is completed by Patient. Response options: Not at all Confident, 1, 2, 3, 4, 5, Completely confident. Reference period not sourced. The result is for clinician review.

CONTEXT

The total is the sum of the 10 item ratings (0-60). No severity band or cut-off is applied. Cutoff not implemented in Lirena.

BOUNDARY

PSEQ is not a diagnosis. A result cannot replace clinician-led assessment, history, examination, or investigation where those are required.

REPORT PRINCIPLE

Show the evidence, the version, and the boundary. Leave diagnosis and formulation with the clinician.

PSEQ response pattern

Synthetic responses for PSEQ. The table uses this instrument's groups and range. It is not a client record and does not reproduce official item wording.

ITEM	GROUP	SYNTHETIC RESPONSE
1	Total	Not at all Confident
2	Total	1
3	Total	2
4	Total	3
5	Total	4
6	Total	5
7	Total	Completely confident
8	Total	Not at all Confident
9	Total	1
10	Total	2

Synthetic PSEQ values only. Not a client record.

Source-led by design

Lirena separates published methodology, product implementation, and clinical approval. A citation does not by itself authorize administration or prove a software scoring rule.

INSTRUMENT

PSEQ captures confidence with function in the presence of pain. It is not a measure of pain intensity and not a personality diagnosis.

ADMINISTRATION

PSEQ is completed by Patient. Response options: Not at all Confident, 1, 2, 3, 4, 5, Completely confident. Reference period not sourced. The result is for clinician review.

CLINICAL BOUNDARY

PSEQ is not a diagnosis. A result cannot replace clinician-led assessment, history, examination, or investigation where those are required.

METHODOLOGY

The total is the sum of the 10 item ratings (0-60). No severity band or cut-off is applied. Cutoff not implemented in Lirena.

SELECTED REFERENCES

1. PSEQ paper

Nicholas MK. The Pain Self-Efficacy Questionnaire: taking pain into account. Eur J Pain. 2007;11(2):153-163.

2. Scoring source Lirena follows

Source URL recorded on the Lirena scoring verification input.