SYNTHETIC EXAMPLE

RCADS

RCADS - synthetic example

CLIENT
John Doe

ADMINISTERED
Example date

RESPONSE MODE
Self-report

TOTAL

106 / 188

DEMO SYNTHETIC



106/188

What this report lays out

- Source instrument: RCADS
- Item count: 47 items
- Response format: None, Mild, Moderate, Severe, Extreme
- Respondent: Self-report

CLINICIAN REVIEW REQUIRED

This document demonstrates Lirena's report design using fictional data. It does not represent a scored RCADS result, cannot establish a diagnosis, and must not be placed in a clinical record.

RCADS does not diagnose an anxiety disorder or depression. Subscale labels are measurement constructs, not confirmed conditions.

A report designed for clinical review

Synthetic RCADS example. The gauges and rows use this instrument's groups and range. They are not a client record.

REVIEW

A young person or parent completes the authorised form. Age, informant, and whether you use the short or long form must be explicit. Results go to a clinician, not a public score page.

CONTEXT

Operational raw-score shell; official RCADS interpretation requires age/sex/informant T-score conversion. The implemented range is 0 to 188. Lirena's implemented cutoff is 65.

BOUNDARY

RCADS does not diagnose an anxiety disorder or depression. Subscale labels are measurement constructs, not confirmed conditions.

REPORT PRINCIPLE

Show the evidence, the version, and the boundary. Leave diagnosis and formulation with the clinician.

RCADS response pattern

Synthetic responses for RCADS. The table uses this instrument's groups and range. It is not a client record and does not reproduce official item wording.

ITEM	GROUP	SYNTHETIC RESPONSE
1	Total	None
2	Total	Mild
3	Total	Moderate
4	Total	Severe
5	Total	Extreme
6	Total	None
7	Total	Mild
8	Total	Moderate
9	Total	Severe
10	Total	Extreme
11	Total	None
12	Total	Mild
13	Total	Moderate
14	Total	Severe
15	Total	Extreme
16	Total	None

Synthetic RCADS values only. Not a client record.

Source-led by design

Lirena separates published methodology, product implementation, and clinical approval. A citation does not by itself authorize administration or prove a software scoring rule.

INSTRUMENT

The Revised Child Anxiety and Depression Scale yields several anxiety-related subscales plus depression. Child and parent versions exist. It is used in many CAMHS-style pathways as a symptom measure, not a diagnostic interview.

ADMINISTRATION

A young person or parent completes the authorised form. Age, informant, and whether you use the short or long form must be explicit. Results go to a clinician, not a public score page.

CLINICAL BOUNDARY

RCADS does not diagnose an anxiety disorder or depression. Subscale labels are measurement constructs, not confirmed conditions.

METHODOLOGY

Operational raw-score shell; official RCADS interpretation requires age/sex/informant T-score conversion. The implemented range is 0 to 188. Lirena's implemented cutoff is 65.

SELECTED REFERENCES

1. RCADS development

Chorpita BF, Yim L, Moffitt C, Umemoto LA, Francis SE. Assessment of symptoms of DSM-IV anxiety and depression in children: a revised child anxiety and depression scale. Behav Res Ther. 2000;38(8):835-855.

2. Developer or publisher page

UCLA RCADS resources