SYNTHETIC EXAMPLE

SPIN

SPIN - synthetic example

CLIENT
John Doe

ADMINISTERED
Example date

RESPONSE MODE
Self-report

TOTAL

34 / 68

DEMO SYNTHETIC



34/68

What this report lays out

- Source instrument: SPIN
- Item count: 17 items
- Response format: Not at all, A little bit, Somewhat, Very much, Extremely
- Respondent: Patient

CLINICIAN REVIEW REQUIRED

This document demonstrates Lirena's report design using fictional data. It does not represent a scored SPIN result, cannot establish a diagnosis, and must not be placed in a clinical record.

SPIN does not diagnose social anxiety disorder. Overlap with depression, panic, and neurodevelopmental presentations still needs clinical judgement.

A report designed for clinical review

Synthetic SPIN example. The gauges and rows use this instrument's groups and range. They are not a client record.

REVIEW

Adults rate 17 items. Results support clinician review. They do not replace a social-anxiety interview.

CONTEXT

The SPIN total is the sum of the 17 items and each subscale is the sum of its items. Connor et al. (2000) reported that a total of 19 distinguished people with and without social phobia. The score does not establish a diagnosis. The implemented range is 0 to 68.

BOUNDARY

SPIN does not diagnose social anxiety disorder. Overlap with depression, panic, and neurodevelopmental presentations still needs clinical judgement.

REPORT PRINCIPLE

Show the evidence, the version, and the boundary. Leave diagnosis and formulation with the clinician.

SPIN response pattern

Synthetic responses for SPIN. The table uses this instrument's groups and range. It is not a client record and does not reproduce official item wording.

ITEM	GROUP	SYNTHETIC RESPONSE
1	Total	Not at all
2	Total	A little bit
3	Total	Somewhat
4	Total	Very much
5	Total	Extremely
6	Total	Not at all
7	Total	A little bit
8	Total	Somewhat
9	Total	Very much
10	Total	Extremely
11	Total	Not at all
12	Total	A little bit
13	Total	Somewhat
14	Total	Very much
15	Total	Extremely
16	Total	Not at all

Synthetic SPIN values only. Not a client record.

Source-led by design

Lirena separates published methodology, product implementation, and clinical approval. A citation does not by itself authorize administration or prove a software scoring rule.

INSTRUMENT

The Social Phobia Inventory asks about fear, avoidance, and physiological symptoms in social situations. It is a brief self-report used for screening or monitoring.

ADMINISTRATION

Adults rate 17 items. Results support clinician review. They do not replace a social-anxiety interview.

CLINICAL BOUNDARY

SPIN does not diagnose social anxiety disorder. Overlap with depression, panic, and neurodevelopmental presentations still needs clinical judgement.

METHODOLOGY

The SPIN total is the sum of the 17 items and each subscale is the sum of its items. Connor et al. (2000) reported that a total of 19 distinguished people with and without social phobia. The score does not establish a diagnosis. The implemented range is 0 to 68.

SELECTED REFERENCES

1. SPIN paper

Connor KM et al. Psychometric properties of the Social Phobia Inventory (SPIN). Br J Psychiatry. 2000;176:379-386.

2. Scoring source Lirena follows

Source URL recorded on the Lirena scoring verification input.