SYNTHETIC EXAMPLE

STOP-Bang

STOP-Bang - synthetic example

CLIENT	ADMINISTERED	RESPONSE MODE
John Doe	**Example date**	**Self-report**

YES COUNT

4 / 8

DEMO SYNTHETIC



What this report lays out

- Source instrument: STOP-Bang
- Item count: 8 items
- Response format: Yes, No
- Respondent: Patient

CLINICIAN REVIEW REQUIRED

This document demonstrates Lirena's report design using fictional data. It does not represent a scored STOP-Bang result, cannot establish a diagnosis, and must not be placed in a clinical record.

STOP-Bang reports the published screen label. It does not diagnose obstructive sleep apnoea.

A report designed for clinical review

Synthetic STOP-Bang example. The gauges and rows use this instrument's groups and range. They are not a client record.

REVIEW

The respondent answers eight yes or no questions. Yes scores 1 and No scores 0. All eight answers are required.

CONTEXT

Lirena uses this rule. Score range: 08. OSA - Low Risk (02), OSA - Intermediate Risk (34), OSA - High Risk (58). OSA - High Risk also applies when questions 14 sum to 2 or more and BMI, neck size, or male sex is Yes. STOP-Bang reports the published screen label. It does not diagnose obstructive sleep apnoea.

BOUNDARY

STOP-Bang reports the published screen label. It does not diagnose obstructive sleep apnoea.

REPORT PRINCIPLE

Show the evidence, the version, and the boundary. Leave diagnosis and formulation with the clinician.

STOP-Bang response pattern

Synthetic responses for STOP-Bang. The table uses this instrument's groups and range. It is not a client record and does not reproduce official item wording.

ITEM	GROUP	SYNTHETIC RESPONSE
1	Yes count	Yes
2	Yes count	No
3	Yes count	Yes
4	Yes count	No
5	Yes count	Yes
6	Yes count	No
7	Yes count	Yes
8	Yes count	No

Synthetic STOP-Bang values only. Not a client record.

Source-led by design

Lirena separates published methodology, product implementation, and clinical approval. A citation does not by itself authorize administration or prove a software scoring rule.

INSTRUMENT

STOP-Bang combines symptom and demographic items used in OSA screening pathways. A screen is not a sleep-study diagnosis and does not by itself decide CPAP or surgery.

ADMINISTRATION

The respondent answers eight yes or no questions. Yes scores 1 and No scores 0. All eight answers are required.

CLINICAL BOUNDARY

STOP-Bang reports the published screen label. It does not diagnose obstructive sleep apnoea.

METHODOLOGY

Lirena uses this rule. Score range: 08. OSA - Low Risk (02), OSA - Intermediate Risk (34), OSA - High Risk (58). OSA - High Risk also applies when questions 14 sum to 2 or more and BMI, neck size, or male sex is Yes. STOP-Bang reports the published screen label. It does not diagnose obstructive sleep apnoea.

SELECTED REFERENCES

1. STOP-Bang paper
Chung F et al. STOP questionnaire. Anesthesiology. 2008;108(5):812-821.

2. Official STOP-Bang site
Toronto Western Hospital. STOP-Bang Questionnaire.

3. Scoring source Lirena follows
Source URL recorded on the Lirena scoring verification input.