SYNTHETIC EXAMPLE

Pain VAS

Pain VAS - synthetic example

CLIENT
John Doe

ADMINISTERED
Example date

RESPONSE MODE
Self-report

PAIN INTENSITY

68 / 100

DEMO SYNTHETIC



68/100

What this report lays out

- Source instrument: Pain VAS
- Item count: 1 item
- Response format: Not sourced
- Respondent: Patient

CLINICIAN REVIEW REQUIRED

This document demonstrates Lirena's report design using fictional data. It does not represent a scored VAS result, cannot establish a diagnosis, and must not be placed in a clinical record.

These ranges describe the mark. They are not a diagnosis and not a treatment recommendation. The recall window is chosen with the form; Hawker records that it is not fixed.

A report designed for clinical review

Synthetic Pain VAS example. The gauges and rows use this instrument's groups and range. They are not a client record.

REVIEW

Pain VAS is completed by Patient. Response format not sourced. Reference period not sourced. The result is for clinician review.

CONTEXT

Lirena uses this rule. Score range: 0100. No pain (04), Mild pain (544), Moderate pain (4574), Severe pain (75100), the ranges described by Jensen, Chen and Brugger (2003). These ranges describe the mark. They are not a diagnosis and not a treatment recommendation. The recall window is chosen with the form; Hawker records that it is not fixed.

BOUNDARY

These ranges describe the mark. They are not a diagnosis and not a treatment recommendation. The recall window is chosen with the form; Hawker records that it is not fixed.

REPORT PRINCIPLE

Show the evidence, the version, and the boundary. Leave diagnosis and formulation with the clinician.

Pain VAS response pattern

Synthetic responses for Pain VAS. The table uses this instrument's groups and range. It is not a client record and does not reproduce official item wording.

ITEM	GROUP	SYNTHETIC RESPONSE
1	Pain intensity	**Synthetic response**

Synthetic Pain VAS values only. Not a client record.

Source-led by design

Lirena separates published methodology, product implementation, and clinical approval. A citation does not by itself authorize administration or prove a software scoring rule.

INSTRUMENT

A pain VAS is a single-intensity rating, not a multidimensional inventory. It does not describe interference, neuropathic features, or function, and it does not diagnose a pain condition.

ADMINISTRATION

Pain VAS is completed by Patient. Response format not sourced. Reference period not sourced. The result is for clinician review.

CLINICAL BOUNDARY

These ranges describe the mark. They are not a diagnosis and not a treatment recommendation. The recall window is chosen with the form; Hawker records that it is not fixed.

METHODOLOGY

Lirena uses this rule. Score range: 0100. No pain (04), Mild pain (544), Moderate pain (4574), Severe pain (75100), the ranges described by Jensen, Chen and Brugger (2003). These ranges describe the mark. They are not a diagnosis and not a treatment recommendation. The recall window is chosen with the form; Hawker records that it is not fixed.

SELECTED REFERENCES

1. Measures of adult pain

Hawker GA et al. Measures of adult pain. Arthritis Care Res. 2011;63(S11):S240-S252.

2. VAS rating ranges

Jensen MP, Chen C, Brugger AM. Interpretation of visual analog scale ratings and change scores. J Pain. 2003;4(7):407-414.

3. Scoring source Lirena follows

Source URL recorded on the Lirena scoring verification input.