SYNTHETIC EXAMPLE

Y-BOCS

Y-BOCS - synthetic example

CLIENT
John Doe

ADMINISTERED
Example date

RESPONSE MODE
Clinician-rated

TOTAL

23 / 40

DEMO SYNTHETIC



23/40

What this report lays out

- Source instrument: Y-BOCS
- Item count: 10 items
- Response format: Varies by item
- Respondent: Clinician

CLINICIAN REVIEW REQUIRED

This document demonstrates Lirena's report design using fictional data. It does not represent a scored Y-BOCS result, cannot establish a diagnosis, and must not be placed in a clinical record.

Y-BOCS does not diagnose OCD. It estimates rated severity. Insight, comorbidity, and functioning still need clinical assessment.

A report designed for clinical review

Synthetic Y-BOCS example. The gauges and rows use this instrument's groups and range. They are not a client record.

REVIEW

A trained clinician interviews the person and records the ten ratings in the clinician completion view. Lirena does not send Y-BOCS to the client.

CONTEXT

The Y-BOCS total is the sum of ten 0-4 ratings, with obsession (items 1-5) and compulsion (items 6-10) subtotals (Goodman et al., 1989). Bands follow the Storch et al. (2015) benchmarks. It is a clinician rating, not a diagnosis. The implemented range is 0 to 40.

BOUNDARY

Y-BOCS does not diagnose OCD. It estimates rated severity. Insight, comorbidity, and functioning still need clinical assessment.

REPORT PRINCIPLE

Show the evidence, the version, and the boundary. Leave diagnosis and formulation with the clinician.

Y-BOCS response pattern

Synthetic responses for Y-BOCS. The table uses this instrument's groups and range. It is not a client record and does not reproduce official item wording.

ITEM	GROUP	SYNTHETIC RESPONSE
1	Total	Synthetic response
2	Total	Synthetic response
3	Total	Synthetic response
4	Total	Synthetic response
5	Total	Synthetic response
6	Total	Synthetic response
7	Total	Synthetic response
8	Total	Synthetic response
9	Total	Synthetic response
10	Total	Synthetic response

Synthetic Y-BOCS values only. Not a client record.

Source-led by design

Lirena separates published methodology, product implementation, and clinical approval. A citation does not by itself authorize administration or prove a software scoring rule.

INSTRUMENT

The Yale-Brown Obsessive Compulsive Scale rates time, interference, distress, resistance, and control for obsessions and compulsions after interview. Symptom checklists used alongside Y-BOCS are not the same as the severity scale.

ADMINISTRATION

A trained clinician interviews the person and records the ten ratings in the clinician completion view. Lirena does not send Y-BOCS to the client.

CLINICAL BOUNDARY

Y-BOCS does not diagnose OCD. It estimates rated severity. Insight, comorbidity, and functioning still need clinical assessment.

METHODOLOGY

The Y-BOCS total is the sum of ten 0-4 ratings, with obsession (items 1-5) and compulsion (items 6-10) subtotals (Goodman et al., 1989). Bands follow the Storch et al. (2015) benchmarks. It is a clinician rating, not a diagnosis. The implemented range is 0 to 40.

SELECTED REFERENCES

1. Original Y-BOCS paper

Goodman WK et al. The Yale-Brown Obsessive Compulsive Scale. Arch Gen Psychiatry. 1989;46(11):1006-1011.

2. Y-BOCS severity benchmarks

Storch EA et al. Defining clinical severity in adults with obsessive-compulsive disorder. Compr Psychiatry. 2015;63:30-35.

3. Scoring source Lirena follows

Source URL recorded on the Lirena scoring verification input.