SYNTHETIC EXAMPLE

YMRS

YMRS - synthetic example

CLIENT
John Doe

ADMINISTERED
Example date

RESPONSE MODE
Clinician-rated

TOTAL

55 / 60

DEMO SYNTHETIC



What this report lays out

- Source instrument: YMRS
- Item count: 11 items
- Response format: Varies by item
- Respondent: Clinician

CLINICIAN REVIEW REQUIRED

This document demonstrates Lirena's report design using fictional data. It does not represent a scored YMRS result, cannot establish a diagnosis, and must not be placed in a clinical record.

A YMRS rating cannot diagnose mania, specify bipolar subtype, or determine treatment. It is one severity estimate inside clinician-led care.

A report designed for clinical review

Synthetic YMRS example. The gauges and rows use this instrument's groups and range. They are not a client record.

REVIEW

A clinician rates items after interview and observation. It should not be converted into an unsupervised questionnaire. Digital capture must keep the rater in the loop.

CONTEXT

YMRS is a clinician-rated severity measure, not a diagnosis. The total is the sum of the 11 ratings; items 5, 6, 8 and 9 carry twice the weight of the others. Young et al. (1978) publish no severity bands. The implemented range is 0 to 60. Cutoff not implemented in Lirena.

BOUNDARY

A YMRS rating cannot diagnose mania, specify bipolar subtype, or determine treatment. It is one severity estimate inside clinician-led care.

REPORT PRINCIPLE

Show the evidence, the version, and the boundary. Leave diagnosis and formulation with the clinician.

YMRS response pattern

Synthetic responses for YMRS. The table uses this instrument's groups and range. It is not a client record and does not reproduce official item wording.

ITEM	GROUP	SYNTHETIC RESPONSE
1	Total	Synthetic response
2	Total	Synthetic response
3	Total	Synthetic response
4	Total	Synthetic response
5	Total	Synthetic response
6	Total	Synthetic response
7	Total	Synthetic response
8	Total	Synthetic response
9	Total	Synthetic response
10	Total	Synthetic response
11	Total	Synthetic response

Synthetic YMRS values only. Not a client record.

Source-led by design

Lirena separates published methodology, product implementation, and clinical approval. A citation does not by itself authorize administration or prove a software scoring rule.

INSTRUMENT

YMRS is used to rate manic symptom severity based on clinician observation and interview. It is common in research and some specialist mood services. It does not diagnose bipolar disorder by itself.

ADMINISTRATION

A clinician rates items after interview and observation. It should not be converted into an unsupervised questionnaire. Digital capture must keep the rater in the loop.

CLINICAL BOUNDARY

A YMRS rating cannot diagnose mania, specify bipolar subtype, or determine treatment. It is one severity estimate inside clinician-led care.

METHODOLOGY

YMRS is a clinician-rated severity measure, not a diagnosis. The total is the sum of the 11 ratings; items 5, 6, 8 and 9 carry twice the weight of the others. Young et al. (1978) publish no severity bands. The implemented range is 0 to 60. Cutoff not implemented in Lirena.

SELECTED REFERENCES

1. Original YMRS paper

Young RC, Biggs JT, Ziegler VE, Meyer DA. A rating scale for mania: reliability, validity and sensitivity. Br J Psychiatry. 1978;133:429-435.

2. Scoring source Lirena follows

Source URL recorded on the Lirena scoring verification input.